



Case Study

SURGICAL MANAGEMENT OF BHAGANDARA BY IFTAK TECHNIQUE ALONG WITH ARAGWADHADI VARTI

Piyush Ranjan Parhi^{1*}, Balendra Singh², Satrupa Nirmal³, Dheeraj Singh Baghel⁴, Subodh Kumar Sahu⁵, Dileep Diwaker⁵

*¹PG Scholar, ²Professor and HOD, ³Lecturer, ⁴Reader, ⁵PG Scholar, Dept. of Shalyatantra, Shri Narayan Prasad Awasthi, Govt. Ayurveda College, Raipur, Chhattisgarh

Article info
Article History:
Received: 30-11-2024
Accepted: 29-12-2024
Published: 15-01-2025

KEYWORDS:

Bhagandara,
Fistula-in-ano,
Ksharasutra,
Aragwadhadi varti.

ABSTRACT

Fistula is a condition in which two epithelial surfaces communicate abnormally. When a communication is established between anal canal and perineal region that condition is called as fistula-in-ano. The track is typically lined with unsightly granulation tissues. The primary cause of this condition is improperly treated crypto glandular infection. Over the advancement of time the *Ksharasutra* is still the best choice among treatment modalities available for Fistula-in-ano because of lesser complications like recurrence and incontinence. Despite all these advantages, there are still certain drawbacks such as discomfort, prolonged periods of anxiety, increased hospital visits, extended treatment duration, and significant post-operative scarring, among others. IFTAK (Interception of Fistulous tract with application of *Ksharasutra*) is a novel advanced *Ksharasutra* technique thus making it more convenient to patient as well as to exclude the drawbacks of conventional method. Further using *Aragwadhadi varti* along with the above said technique lead to better healing of the leftover portion of the track. Here in this case, IFTAK is done under Local anaesthesia. *Kshara sutra* was changed upto 4 weeks and *Aragwadhadi varti* applied for 10 times. The fistula healed completely in 2 months. This study revealed an early complete remission of the fistulous tracts by IFTAK method and *Aragwadhadi varti*, When compared with the conventional *Kshara sutra* method. According to the length, the conventional method takes 15-20 weeks whereas in this study it took much less time. And during followup no recurrence was noted. Therefore, the IFTAK technique combined with *Aragwadhadi varti* was deemed highly effective due to its time-saving properties and reduced scar and fibrosis.

INTRODUCTION


The untreated or improperly treated cryptoglandular infection is the most common cause of fistula in ano or *Bhagandara*.^[1] In this case study administration of IFTAK (Interception of Fistulous track with application of *Ksharsutra*) technique was done along with application of *Aragwadhadi Varti* in treating a complex intersphincteric fistula in Ano successfully. Although the disease is not that much life

threatening but produces severe inconvenience in routine life because of pus discharge and pain.

By looking into the severity and chronicity of disease The Father of surgery Acharya Sushruta included *Bhagandara* in *Asta Mahagada*.^[2] There are numbers of treatment modalities available for the management of fistula in ano. Modern surgical management includes fistulotomy, fistulectomy, Seton placing, ligation of intersphincteric fistula tract (LIFT), fibrin glues, advancement flaps, and expanded adipose derived stem cells (ASCs).^[3] Acharya Sushruta has also described different therapeutic measures for the management *Bhagandara* as in terms of various oral medications, local applications, surgical procedures and para-surgical intervention. Acharya Charaka has mentioned it under *Swayathu chikitsa adhyaya* and *Kshara sutra* ligation as its treatment^[4] Presently

Access this article online	
Quick Response Code	https://doi.org/10.47070/ayushdhara.v11i6.1867
	Published by Mahadev Publications (Regd.) publication licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0)

Ksharasutra therapy is found most approaching and attractive treatment modality among para-surgical procedure for fistula-in-ano. Over the advancement of time the *Ksharasutra* is still the best choice among treatment modalities available for Fistula-in-ano because of lesser complications like recurrence and incontinence. But with all these benefits, still there are some consequences suffered by practitioners while practicing *Ksharasutra* therapy in relation to patients i.e. it causes discomfort, long anxiety period, number of hospital visits and longer duration of treatment, big post-operative scar etc. IFTAK (Interception of Fistulous tract with application of *Ksharasutra*) is an innovative *Ksharasutra* technique rooted in this concept, thereby enhancing convenience for patients and avoiding the limitations of traditional methods. [5] There are a number of choices in *Ksharasutra* like *Apamarga* based, papaya based, *Guggulu* based or *Palasha* based. But in this case we need to use *Apamarga Ksharasutra* because it has terrific result in cutting and healing the fistulous track. [6] Thus, cases of fistula treated with this method can produce positive



Clinical Findings

On examination perianal skin was normal. An external opening was present at 3 O'clock approximately 7 cm away from anal verge. On digital rectal examination, sphincter tone was normal tender dimpling or buttonhole-like opening noted at 3 O'clock at a distance of 1.5 cm inner to OS. On probing it was found that the probe goes straight towards anal verge in a curvilinear manner. A single complete fistulous tract of about 7.5cm in length was found. Rest of the examinations were found to be normal and laboratory investigations were also found within normal limits.

results. Additionally, utilizing *Aragwadhadi varti* alongside the aforementioned technique might enhance the healing of the remaining track portion and reduce the likelihood of subsequent sinus recurrences. Because the indurated portion of the fistulous tract also needs to be healed properly along with the area near anal verge. *Aragwadhadi varti*, as mentioned by Acharya Sushruta in the *Bhagandara roga adhikara*, may offer improved outcomes in reducing fibrosis of the fistulous tract and facilitate quicker healing compared to the standard treatment.

Presenting Complaints and Medical History

A 31 year old male patient came to Shalya OPD of Shri Narayan Prasad Awasthi Govt. Ayurveda College, Raipur having Complaints of recurrent gluteal abscess since last 2 years along with the complaints of Intermittent fever and pus discharge from anal verge. Patient was not having any Comorbidities. Patient got operated with laser around 8 months prior. Endoanal X-Ray fistulogram reports are attached below.



Treatment

After obtaining written consent and completing necessary pre operative procedures, the patient was placed in lithotomy position. After painting and draping, probing was done to ascertain the finding. After confirming the diagnosis, a small vertical incision was made at 3 O'clock position at a distance of 1 cm near from anal verge and Interception of fistulous tract was done in intersphincteric space. Then a metallic probe was introduced through the window and taken out from the internal opening and *Kshara sutra* ligation was done in the proximal tract. The *Kshara sutra* was ligated loosely on initial days in order to facilitate proper drainage of pus. The residual distal tract was also ligated with *Ksharasutra* in order to keep the tract patent (Fig.7). After achieving haemostasis dressing

and packing done with *Jatyadi taila* and patient then shifted to male surgical ward. Patient was advised for regular hot sitz bath from post operative day 2. Patient was prescribed with *Triphala guggulu* 2 tab BD after food, Syp. *Abhayarista* 15 ml BD with equal amount of water after food, *Triphala churna* 5 gm HS with luke warm water after food.

On dressing from post operative day 2, *Ksharasutra* was removed from distal tract and antiseptic dressing done with *Jatyadi Taila* and *Aragwadhadi varti* inserted in distal tract on every alternate days till the signs of *Shuddha vrana* appeared.

Aragwadhadi Varti

In Sushruta Samhita, *Aragwadhadi varti* is described in Treatment of *Bhagandara*.^[7] *Aragwadhadi varti* has properties of *Shodhana* i.e., Cleanses the tissue by scraping out unhealthy tissue and *Ropana* i.e., Accelerates healing process by clearing debris tissue.^[8] Powdered *Haridra* (*Curcuma longa*) and *Tagara* (*Valeriana wallichii*) taken in equal quantities and mixed in a well-balanced manner, followed by addition

of *Aragwadha* (*Cassia fistula*) *Majja*, *Ghritha*, and honey, all in equal quantities (Fig.3). The wicks were then prepared by hand, measuring 2 to 3 inches in length, and dried in the sun (Fig.4). Once the wicks dried and hardened, they were sterilized in UV chambers and then placed in air-tight containers. *Aragwadhadi varti* was prepared in the Department of Rasa Shastra and Bhaishajya kalpana of Shri N.P.A. Govt. Ayurveda College Raipur (C.G) according to classical reference.

The ingredients of formulation *Aragwadhadi varti* were collected from local fields and purchased from the market (Fig.1 & 2) and were authenticated by help of the Department of Dravyaguna, Govt. Ayurveda college Raipur (C.G). Moreover, further authentication was done from the Central research facility, KLE Academy of higher Education and Research, Karnataka. *Aragwadhadi varti* was sent to the Drug Testing Laboratory and Research Centre, Raipur (C.G) / NABL accredited lab for analysis of phytochemical and physiochemical constituents.



Figure 1



Figure 2



Figure 3




Figure 4

Followup and Outcomes

Weekly follow up advised for *Ksharsutra* changing. *Ksharsutra* was changed three times after the first *Ksharsutra* was placed. The pus discharge was fluent in the first week from the artificially made window, gradually reduced and completely

disappeared after two weeks (Fig. 7). Pain was also moderate in the first week and later on gradually relieved. The discharge from the external opening was also reduced gradually in 6-7 days and totally dried up in three weeks. (Fig. 8) Then the *Ksharasutra* was

ligated tightly and Cut through was done (Fig. 9) when discharge completely diminished from the artificial made window i.e., after three weeks of first *Ksharasutra* placed and complete healing was achieved in 15 days after cut through of proximal tract. Patient was advised for application of *Jatyadi taila*. The fistulous tract healed completely by 6th week with minimal scar and without any fibrosis (Fig. 10). There was no complication seen during and after treatment and patient got free from all the symptoms. After 12 months of follow up, no recurrence is noted, the patient was cured completely.

Probable Mode of Action of *Aragwadhadi Varti*

Traditional treatments for wound healing include honey, *Curcuma longa*, *Valerian wallichii*, and *Cassia fistula*. *Cassia fistula*'s anti-inflammatory, antibacterial, antifungal, and antioxidant qualities aid in wound closure, tissue regeneration, and enhanced healing. [9] *Curcuma longa* contains antibacterial, antioxidant, and anti-inflammatory qualities[10], while *Valerian wallichii* possesses analgesic and anti-

inflammatory qualities. [11] As a natural antimicrobial, honey promotes quicker healing and reduced scarring by lowering inflammation, encouraging tissue development, and aiding autolytic debridement.[12] *Ghrita* is a traditional Ayurvedic medication that decreases inflammation and speeds up wound contraction. Also it has been found that *Aragwadhadi varti* has a pH of 4.62, which is slightly acidic. Because it encourages quicker re-epithelialization, collagen deposition, and wound closure, a slightly acidic environment is optimal for wound healing. [13] It also prevents the formation of dangerous microorganisms. Acidification promotes the migration of epithelial cells, increases oxygen release, and activates fibroblasts. By preventing the development of fibroblasts into myofibroblasts, which are in charge of collagen deposition and scarring, it may be able to lessen fibrosis.[14] Collagen synthesis and fibrosis development can be inhibited by lowering pH levels in cell culture conditions. [15]



Figure 5



Figure 6



Figure 7



Figure 8



Figure 9



Figure 10

DISCUSSION

Ksharasutra therapy is the most successful treatment modality for fistula in ano. *Ksharasutra* therapy has high success rate[16] and least recurrence rate (3.33%)[17]. It is a highly simple day care option and an affordable treatment with a lower complication rate compared to traditional treatment methods that

involve hospitalization, regional or general anesthesia, and frequent post-operative care. These surgical procedures carry a notable risk of recurrence (0.7–26.5%) and a considerable risk of reduced continence (5–40%).[18] While *Ksharasutra* therapy is a preferred treatment for fistula in ano due to its various benefits,

it does come with certain drawbacks, including discomfort, pain after the procedure, multiple hospital visits, extended treatment duration, significant post-operative scarring, and post-treatment fibrosis, resulting in low compliance and reduced acceptability among many patients. IFTAK (Interception of Fistulous tract and application of *Ksharsutra*) technique seems to overcome the limitations and consequences of conventional methods. Duration of therapy was less by shortening the length of the track and taking care of crypto glandular infection where there was no need to treat residual curved track. Pain was significantly reduced because of less exposure of tissues after interception which is from internal opening where as in conventional method whole track was exposed along the axis during the *Ksharsutra* change which increases the pain and burning sensation because of more tissue exposure.

In this case study, the external opening was approximately 7 cm from the anal verge at the 3 o'clock position, and interception occurred at the 3 o'clock position about 1 cm from the anal verge, thus decreasing the length of the tract. Consequently, the IFTAK technique shortened the total healing time and significantly decreased pain while leaving minimal scarring.^[19] *Aragwadhadi varti* purified the leftover tract and the sinus tract due to its *Shodhana* and *Ropana* properties. The patient was fully healed within two months, and no recurrence was observed during the three-month follow-up.

Tablet *Triphala Guggulu* were used to counter inflammation, pain and to prevent infection. *Triphala Guggulu* tablets were utilized to alleviate inflammation, relieve pain, and guard against infection. *Triphala guggulu* has demonstrated antimicrobial properties.^[20] Thus, it may be useful in prevention of infection and promote wound healing also. *Triphala* has antibacterial action against a variety of Gram-positive and Gram-negative bacteria and immunomodulatory effect.^[21] And as per Ayurveda, *Triphala* exhibits *Anulomana* properties, which helps balance *Apana Vata* and promotes smooth bowel movements. The local use of *Jatyadi Taila* is said to notably enhance the levels of protein, hydroxyproline, and hexosamine in granulation tissue, facilitating quicker healing.^[22] So, IFTAK is a sophisticated and efficient method in the area of fistula that features significant innovation and numerous advantages. Additionally, *Aragwadhadi varti* functions like the cherry atop a cake when it comes to cleaning the fistulous tract, enhancing wound healing, and preventing fibrosis following complete recovery.

CONCLUSION

Therefore, the research concluded that IFTAK, in conjunction with the use of *Aragwadhadi Varti*, is a secure, efficient, and progressive method that reduces post-operative duration while improving mild post-procedural pain and minimizing scar marks.

REFERENCES

1. Kumar A, Bilyan A. IFTAK an innovative technique in fistula in ano-A case study. *Ayur pub* 2018; 2:771e5.
2. Sushruta Samhita, Part-I, Ayurveda Tatwa Sandeepika Hindi Commentary, Edited by Kaviraj Ambika Dutta Shastri, Sutrastana- Avarniya Adhyaya 33/4, Chaukhamba Sanskrit Sansthan, Varanasi, Edition-13, 2002, Page no. 126.
3. Limura E, Giordano P. Modern management of anal fistula. *World J Gastroenterol* 2015;21(1):12e20. <https://doi.org/10.3748/wjg.v21.i1.12>.
4. Charaka Samhita, part 2, uttarardha, Vidyotini Teeka Hindi commentary, edited by Kashinath pandey shastri, chikitsa sthana – swayathu chikitsa 12/97, Chaukhamba Sanskrit sansthan, Varanasi, Edition 13, 2021, Page no – 342.
5. Diwan S, Kumar P, Gupta SJ. IFTAK-An advanced technique of Kshara Sutrathrapy in management of complex Fistula-In-Ano-A Case Study. *J Ayurveda Integrated Med Sci* 2019;3(6):181e4
6. Kumar A. Current trends in the usage of ksharakarma (alkaline therapy) and ksharasutra (alkaline seton) for managing bhagandhara (fistula in ano). *J. res. tradit. med* 2018;4(2):70e7.
7. Sushruta, Sushruta Samhita, Part-I, Ayurveda Tatwa Sandeepika Hindi Commentary, Edited by Kaviraj Ambika Dutta Shastri, chikitsa stana-Bhagandara chikitsa Adhyaya 8/30, Chaukhamba Sanskrit Sansthan, Varanasi, Edition-13, 2002, Page no. 60.
8. Sushruta, Sushruta Samhita, Part-I, Ayurveda Tatwa Sandeepika Hindi Commentary, Edited by Kaviraj Ambika Dutta Shastri, chikitsa stana-Bhagandara chikitsa Adhyaya 8/30, Chaukhamba Sanskrit Sansthan, Varanasi, Edition-13, 2002, Page no. 60.
9. Senthil Kumar M, Sripriya R, Vijaya Raghavan H, Sehgal PK. Wound healing potential of Cassia fistula on infected albino rat model. *J Surg Res.* 2006 Apr; 131(2):283-9. doi: 10.1016/j.jss.2005.08.025. Epub 2005 Oct 20. PMID: 16242721.
10. Tejada S, Manayi A, Daglia M, Nabavi SF, Sureda A, Hajheydari Z, Gortzi O, Pazoki-Toroudi H, Nabavi SM. Wound Healing Effects of Curcumin: A Short Review. *Curr Pharm Biotechnol.* 2016;17(11):

- 1002-7. doi: 10.2174/13892010176661607211 23109. PMID: 27640646.
11. Li J, Li X, Wang C, Zhang M, Ye M, Wang Q. The potential of Valeriana as a traditional Chinese medicine: traditional clinical applications, bioactivities, and phytochemistry. *Front Pharmacol*. 2022 Sep 21;13:973138. doi: 10.3389/fphar.2022.973138. PMID: 36210806; PMCID: PMC9534556.
 12. Ahmad Oryan, Esmat Alemzadeh, Ali Moshiri, Biological properties and therapeutic activities of honey in wound healing: A narrative review and meta-analysis, *Journal of Tissue Viability*, Volume 25, Issue 2, 2016, Pages 98-118, <https://doi.org/10.1016/j.jtv.2015.12.002>. (<https://www.sciencedirect.com/science/article/pii/S0965206X15000972>)
 13. Sim P, Strudwick XL, Song Y, Cowin AJ, Garg S. Influence of Acidic pH on Wound Healing In Vivo: A Novel Perspective for Wound Treatment. *Int J Mol Sci*. 2022 Nov 7;23(21):13655. doi: 10.3390/ijms232113655. PMID: 36362441; PMCID: PMC9658872.
 14. Sim P, Song Y, Yang GN, Cowin AJ, Garg S. In Vitro Wound Healing Properties of Novel Acidic Treatment Regimen in Enhancing Metabolic Activity and Migration of Skin Cells. *Int J Mol Sci*. 2022 Jun 28;23(13):7188. doi: 10.3390/ijms23137188. PMID: 35806191; PMCID: PMC9266998.
 15. Sim P, Song Y, Yang GN, Cowin AJ, Garg S. In Vitro Wound Healing Properties of Novel Acidic Treatment Regimen in Enhancing Metabolic Activity and Migration of Skin Cells. *Int J Mol Sci*. 2022 Jun 28;23(13):7188. doi: 10.3390/ijms23137188. PMID: 35806191; PMCID: PMC9266998.
 16. Pankaj S, Manoranjan S. Efficacy of Ksharsutra (medicated seton), therapy in the management of fistula-in-ano. *World J Colorectal Surg* 2010;2(2) (Art. 6:01e10).
 17. Panigrahi HK, Rani R, Padhi MM, Lavekar GS. Clinical evaluation of Ksharasutra therapy in the management of Bhagandara (fistula-in-ano)da prospective study. *Anc Sci Life* 2009;28(3):29e35 [PMC free article] [PubMed].
 18. Dutta G, Bain J, Ray AK, Dey S, Das N, Das B. Comparing Ksharasutra (Ayurvedic Seton) and open fistulotomy in the management of fistula-in-ano. *J NaSci Biol Med* 2015;6(2):406e10. <https://doi.org/10.4103/0976-9668.160022>. PMID: 26283840; PMCID: PMC4518420.
 19. Sahu M. published by. In: A manual on fistula in ano and kshara sutra therapy. 1st ed. NRC, Deptt. Of ShalyaTantra, IMS, BHU; 2015.
 20. Mhaiskar Bhushan D, Bharat Chouragad Bari. Management of non-healing infected wound by external application of and Haridradya Taila Triphala Guggulu. *Joinsysmed* 2017;5(2):130e4.
 21. Tarasiuk A, Mosinska P, Fichna J. Triphala: current applications and new perspectives on the treatment of functional gastrointestinal disorders. *Chin Med* 2018;13:39. <https://doi.org/10.1186/s13020-018-0197-6>. PMID: 30034512; PMCID: PMC6052535.
 22. Shailajan S1, Menon S, Pednekar S, Singh A. Wound healing efficacy of Jatyadi Taila: in vivo evaluation in rat using excision wound model. *J Ethnopharmacol* 2011;138(1):99e104. <https://doi.org/10.1016/j.jep.2011.08.050>. Epub 2011 Aug 30.

Cite this article as:

Piyush Ranjan Parhi, Balendra Singh, Satrupa Nirmal, Dheeraj Singh Baghel, Subodh Kumar Sahu, Dileep Diwaker. Surgical Management of Bhagandara by IFTAK Technique Along with Aragwadhadi Varti. *AYUSHDHARA*, 2024;11(6):355-360. <https://doi.org/10.47070/ayushdhara.v11i6.1867>

Source of support: Nil, Conflict of interest: None Declared

***Address for correspondence**

Dr. Piyush Ranjan Parhi

PG Scholar,
Dept. of Shalyatantra,
Shri Narayan Prasad Awasthi,
Govt. Ayurveda College, Raipur,
Chhattisgarh.

Email:

parhi.piyushranjan@gmail.com

Disclaimer: AYUSHDHARA is solely owned by Mahadev Publications - A non-profit publications, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. AYUSHDHARA cannot accept any responsibility or liability for the articles content which are published. The views expressed in articles by our contributing authors are not necessarily those of AYUSHDHARA editor or editorial board members.