



Review Article

THERAPEUTIC EFFICACY AND PHYTOCHEMICAL INSIGHTS INTO THE CLASSICAL UNANI FORMULATION *NUSKHA-E-KHALAL-E-SHIKAM*

Uzma Viqar^{1*}, Sadaf Joweria², Mubasshera Mohd. Irfan³, Mohd. Noman Taaha⁴, Younis Iftikhar Munshi⁵

¹Professor, ³PG Scholar, PG Dept. of Ilmul Advia (Unani Pharmacology), National Research Institute of Unani Medicine for Skin Disorder, Hyderabad, Telangana.

²Assistant Professor Dept. of Ilmul Advia (Unani Pharmacology), Markaz Unani Medical College & Hospital, Kozhikode, Kerala.

⁴Assistant Professor, Dept. of Ilmul Advia (Unani Pharmacology), Deoband Unani Medical College and Research Center, Saharanpur, UP.

⁵Deputy Director General, Central Council for Research in Unani Medicine (CCRUM), Ministry of Ayush, Govt. of India, New Delhi, India.

Article info

Article History:

Received: 18-03-2026

Accepted: 22-04-2026

Published: 06-05-2026

KEYWORDS:

Nuskha-e-Khalal-e-Shikam, Mushtarak-un-nafa'a advia, Humma.


ABSTRACT

In the Unani System of Medicine, the concept of *Mushtarak-un-nafa' advia* has existed since ancient times. Certain selected *Advia* both single and compound drugs are outlined under this heading. *Mushtarak-un-nafa' advia* refers to medications that are frequently administered for the majority of illnesses; these medications can be administered to patients without waiting for a proven diagnosis, even if they do not successfully treat the illness or even make it worse. *Mushtarak-un-nafa' advia* is exemplified by *Nuskha-e-Khalal-e-Shikam* (NKS) and *Mulayyin* medicines. NKS composed of five ingredients, i.e., *Gul-e-Banafsha*, *Beekh Kasni*, *Badiyaan*, *Gaozaban* and *Maveez Munaqqa*. NKS is used as *Munjiz* and *Mushil* and is very efficacious in *Ufooni humma* (*Safrawi* or *Balghami*). NKS has traditionally been utilized in situations where a patient's illness is difficult to diagnose. NKS can be modify according to the condition of the patient. Gut dysbiosis is implicated in the pathogenesis of numerous inflammatory diseases. *Nuskha-e-Khalal-e-Shikam* (NKS) ingredients are rich in potent phytochemicals with a range of pharmacological actions, which together produce its compound therapeutic benefits across different conditions. However, more research is needed to assess NKS's effectiveness as a compound formulation.

INTRODUCTION

In India and other areas of the world, the Unani System of Medicine (USM) is among the oldest medical practices. The Unani System of Medicine (USM) originated in Greece and was refined into a sophisticated medical discipline by Arabs, building on the teachings of Hippocrates and Galen. USM advocates the Hippocratic (*Buqrat*) theory of four humours

(*Akhlāt*) i.e., *Dam* (Blood), *Balgham* (Phlegm), *Safra* (Bile), and *Sawda* (Black bile) which have specific but different temperament (*Mizāj*) [1,3-5]. The majority of the diseases arise from a qualitative or quantitative impairment of temperament, humor, or both. Unani medicine attempts to address health-related problems with the aid of natural ingredients and has its own philosophical and theoretical framework. Humoural imbalance is often the root cause of diseases in USM. According to Unani physicians, the human body is composed of seven basic principles (*Umoor-e-Tabbiya*)- *Arkān* (element), *Mizāj* (temperament), *Akhlāt* (humours), *A'ḍā'* (organs), *Arwah* (vital force), *Quwā* (faculties) and *Af'āl* (functions). Disease is caused by the simple lack of any one of the components, endangering life itself. The primary goal

Access this article online	
Quick Response Code	
	https://doi.org/10.47070/ayushdhara.v13i2.2557
Published by Mahadev Publications (Regd.) publication licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0)	

of a Unani physician is to help the body's faculties in order to restore equilibrium. *Ṭabī'a-e-Mudabbira-i-Badan* (Medicatrix naturae) [1-3].

Unani physicians also introduced the idea of using *Mushtarak-un-nafa' advia* (*Da'eef ul-a'mal*) for patients with ambiguous conditions, despite the existence of numerous well-known and recognized philosophical and theoretical concepts. where a precise diagnosis of the disease is difficult [2,4]. Even with today's tremendous advancements in medicine, there are still several conditions of the patients where to reach the confirmative diagnosis is very difficult even after the proper screening of the patient. Pyrexia of unknown origin (PUO) is a prime example- it's one of those tough medical puzzles that really challenges doctors. According to Petersdorf and Beeson, PUO is characterized by a temperature that is higher than 38.3 degrees Celsius, develops over a period of at least three weeks, with no diagnosis reached after one week of inpatient investigation. While addressing such issues, it is imperative to utilize all available clinical and investigative techniques while keeping the local epidemiological conditions in mind. Infections continue to be the most significant cause of PUO, accounting for roughly 20–40% of cases, despite some variance in the range of diseases observed in other series addressing PUO [6,7].

The definition of pyrexia in itself is complex as there is no agreed consensus. The inaccuracy of peripheral thermometers in determining body core temperature further complicates this. Pyrexia has a variety of causes, which lead to various definitions. Fever is typically defined as a temperature higher than 38.3 °C during infection. In the post-resuscitation management of cardiac arrest, a threshold is 37.6°C; in stroke, thresholds of 37.2, 37.5 and 38°C are used. A core temperature of less than 36°C is typically indicative of hypothermia, regardless of the clinical context [8].


The Unani System of Medicine provides a very detailed description of the fever (*Humma*). The various types of fever along with their pathophysiology, clinical features, and management strategies are explained with exceptional clarity.

MATERIALS AND METHODS

Literature for this review was systematically searched from classical Unani texts, including

Kulliyat-e-Asri, Kulliyat-e-Nafeesi, Kulliyat-e-Qanoon, Sharah Asbab, Hummiyat-e-Qanoon, Khazāin al-Adwiyah, Muḥīṭ-i A'zam, and Makhzan-ul-Mufradāt (al-marūf bi-Khawāṣṣ al-Adwiyah). Additionally, the Unani Pharmacopoeia of India and Standard Unani Medical Terminology were referred to ensure accurate description of Unani terminologies. English-language books, research articles, review papers, the Glossary of Indian Medicinal Plants, and publications from journals (PubMed, Google Scholar, and ResearchGate) were also referred to compile evidence-based insights on the formulation's composition, pharmacology, and therapeutic efficacy.

Classification of fever (*Ḥummā*) [9]



The methods of treatment according to USM is divided into four different parts namely Dietotherapy (*Ilāj bi'l Ghidhā*), Regimental therapy (*Ilāj bi'l Tadbīr*), Surgery (*Ilāj bi'l Yad*) and Pharmacotherapy (*Ilāj bi'l Dawā*) [2,3]. In Unani medicine, therapeutic agents are utilized to restore the body's humoral balance and restore health. Maintaining the humoral balance is also thought to depend on a balanced diet and healthy digestion. Pharmacotherapy involves the use of both single and complex medications. Pharmacotherapy deals with the use of naturally occurring drugs, primarily herbal, though it also includes medications derived from animals and minerals. Since USM places a strong emphasis on each person's unique temperament, medications are administered based on the patient's temperament, thus accelerating the process of recovery and also eliminating the risk of drug reaction. Although single medications are often preferred in USM, compound formulations are also utilized to treat a variety of diseases.

Types of <i>Ḥummā</i> [9]	Treatment
<i>Ḥummā al-Yawm</i>	Can be cure with various modalities
<i>Ḥummā Diqqiyya</i>	Specific medication is needed
<i>Ḥummā 'Ufūniyya/Khilṭi</i> (on the basis of predominant temperament)	<ul style="list-style-type: none"> ▪ <i>Mushtarak tadābīr</i> (<i>Talliyun, Ta'reeq, Qai</i> etc.) ▪ <i>Mushtarak nafa' adwia</i> (<i>Nuskha-khalal-e-shikam</i>) ▪ Specific treatment after confirmed diagnosis.

In medical conditions where the clear diagnosis of the disease is difficult, Unani Physician used the *Mushtarak-un-nafa' adwia (Da'eef ul-a'mal)* which if not effectively cure the disease, nor even worsen the condition. Keeping in view of this problem Unani physician formulates a unique formulation for all types of fever, irrespective of its diagnosis. *Nuskha e khalal e shikam (NKS)* can be used in all types of *Ḥummā 'Ufūniyya* as a provisional diagnosis, till the confirmation of final diagnosis, without hesitation, it has good results [2-4,9-10]. This nuskha NKS can also be

given to patients of Gastrointestinal related problems. Similarly, if the patient suffers from coryza, cough or any respiratory disorder of which is difficult to reach the confirmed diagnosis, this Nuskha NKS can be used effectively and if necessary, the drug specific for coryza like *Aslussos, Tukhm-e-Khatmi* etc can be added to NKS [9,10].

According to confirmative diagnosis, this NKS can be modify later according to the related associated medical conditions of the patient.

Ingredients of Nuskha-e-khalal-e-Shikam is as follows [2-4,9-10]

S.No	Unani Name	Botanical Name [11]	Parts Used	Mizaj	Quantity [4,9,11]
1.	<i>Gul-e- Banafsha</i>	<i>Viola odorata</i> L.	Flower	Cold- 1 Moist-1 [12-14]	7 gm
2.	<i>Beekh kasni</i>	<i>Cichorium intybus</i> L.	Root	Cold- 1 Moist- 1[12-14]	7 gm
3.	<i>Badyan</i>	<i>Foeniculum vulgare</i> Mill	Seeds	Hot- 2 Dry- 2[13-14]	7gm
4.	<i>Gaozaban</i>	<i>Borago officinalis</i> L.	Leaves	Hot- 1 Moist- 1[12,14]	5 gm
5.	<i>Maweez munaqqa</i>	<i>Vitis vinifera</i> L.	Fruit	Hot-1 Moist-1[12-15]	9 no.

Method of preparation and administration

All the dry ingredients (mentioned above in the table) are soaked in warm water overnight and boiled next morning till only one-fourth volume of water remains. The ingredients of the decoction is now rubbed with the hands, then filtered through a sieve. The lukewarm decoction is administered orally in the morning hours to the patient along with *Khameera Banafsha* 40ml [4,9,10].

NKS can be modify according to condition of patient^[10]

➤ If along with fever presents

- Sore throat (*Dard-e-Halaq*) then add *Barg-e-Toot* in NKS.
- Coryza (*Nazla*) then add *Tukhm-e-Khatmi* and *Tukhm-e-Khubbazi* in NKS.
- Cough (*Khansi*) then add *Aslussos* in NKS.
- Spleenomegaly (*Waram-e-Tihāl*) then add *Anjeer Zard* in NKS.

○ Dysentery and *Kharash-e-Ama'a* then add *Luab-e-Resha-e-Khatmi* in prepared decoction in morning.

○ *Fasād-e-Khoon* or Skin diseases then add *Shahtara* and *Chiraita*.

○ Amenorrhoea (*Ihtibās-e-Haiz*) then add *Tukhm-e-Khayarain* and *Tukhm-e-Kasni* in NKS.

○ Pallor (*Yaraqān*) then add *Tukhm-e-Khyarain* in NKS and add *Aab-e-Barg-e-Turb* in decoction prepared in the morning.

➤ After the seven days use of this NKS, still the fever persists then add *Khaksi Pasheeda* in NKS.

➤ If *Tahlil-e-Ruṭūbat* and *Talṭif-e-Mādda* is prime concern then add *Beekh-e-Izkhar* and *Beekh-e-Kibr* in the NKS.

➤ To relieve the constipation, add *Gul-e-Surkh* in NKS when coryza is not present, otherwise add *Sana makki* or *Rewand chini* in NKS.

Brief description of Ingredients

S.No	Drug	Chemical Constituents	Pharmacological Action	Pharmacological Studies
1.	<i>Gul-e-Banafsha</i> Arabic: <i>Banafsaj</i> ,	Flavonoids, tannins, alkaloid, glycoside, saponins, different essential oils (methy ester, salicylic acid,	<i>Mu'addil</i> , <i>Mulattif</i> (demulcent), <i>Mu'arriq</i> (diaphoretic), <i>Mulayyin-i-Shikam</i>	Anti-inflammatory activity [21] Antioxidant activity and radical scavenging activity [22] Antimicrobial activity/antibacterial

	English: Violet, Sweet violet ^[15]	anthocyanin, gamma sitosterol, phytol, octadecanoid acid), cycloviolacin, eugenol, violin, rutin, tocopherol. ^[16-20,22]	(Laxative), <i>Mulayyin-i-Halaq-wa-Seenah</i> , <i>Musakkin -e-Dam</i> , <i>Murattib</i> (humactant), <i>Munawwim</i> (hypnotic), <i>Mushil</i> (purgative), <i>Mushil-i-Safra'</i> (Cholagogue), <i>Muzliq</i> , <i>Dāfi'-i-'Uṭāsh</i> , <i>Muḥallil</i> (resolvent) ^[12-14,18]	activity ^[22] Immunological and cytogenic effect ^[23] Hepato-protective ^[23] Expectorant ^[24] Laxative ^[25] Diuretic activity ^[25] Antipyretic activity ^[26] Anti-hypertensive and dyslipidemic activity ^[27] Anti-tubercular activity ^[28] Hypnotic activity ^[29] Pancreatic lipase inhibitors activity ^[30]
2.	<i>Beekh kasni</i> Arabic: <i>Hindyba</i> English: Endive ^[16,18]	Phytosterols, lactones, flavonoids, triterpenoids, terpene, coumarins (including cichorin), tannins, vitamins, pectins. sesquiterpenes lactones like sonchusides A and C, and, cytokinin, crepidiase B, cichoriolide A, B and C, chlorogenic, neochlorogenic, isochlorogenic, lactucin, caffeic & chicoric acids. Series of glucofructosans. ^[16,18,31,32]	<i>Mufattiḥ-i-Sudur</i> (Deobstruent), <i>Muddir-i-Bawl</i> (diuretic), <i>Muṣaffi-i-Dam</i> (blood purifier), Reduces the <i>Ḥarārat</i> of <i>Dam</i> and <i>Ṣafra'</i> , <i>Muqawwī-i-Kabid</i> (hepatotonic), <i>Muqawwī-i-Mi'da</i> (stomachic), <i>Muskkin-i-'Uṭāsh</i> (Thirst reliever), <i>Musakkin</i> (analgesic), <i>Muḥallil</i> (resolvent) ^[12-14,18,33]	Antioxidant activity ^[34,37] Hepatoprotective activity ^[34,35] Antiulcer activity ^[36,37] Antibacterial activity ^[38] Antidiabetic activity ^[39] Antifungal activity ^[40] Anticancer activity ^[41] Anti-convulsant activity ^[42] Immunomodulator activity ^[43] Analgesic and Sedative activity ^[44] Cardioprotective Activity ^[45]
3.	<i>Badyan</i> English- Fennel Arabic: <i>Rajiyanaaj</i> Urdu: Saunf ^[15,16,18]	Volatile oil (anethole, fenchone, methylchavicol, camphene, terpinene, linalool, estragole, p-cymene, thujene), Fixed oil (petroselinic acid), Cinnamic acid derivatves (Hydroxycinnamic acids, cynarin), Flavonoids glycosides (3-glucuronides of kaempferol, Quercetin), Coumarins	<i>Mufattiḥ-i-Sudur</i> (deobstruent), <i>Muqawwī-i-Mi'da</i> (stomachic), <i>Muddir-i-Bawl</i> (diuretic), <i>Mudirr-i-Ḥayḍ</i> (emmenagogue), <i>Muqawwī-e-Başar</i> (vision improving), <i>Mundij-i-Balgham wa Sawdā'</i> (concoctive of phlegm and black bile), <i>Muwallid-i-Laban</i> (galactopoietic), <i>Mujaffif</i> (Desicant), <i>Muwallid-i-Manī</i> (spermatogenic),	Antifungal activity ^[48] Antioxidant activity ^[49] Anti-inflammatory ^[49] Analgesic ^[49] Antidiabetic activity ^[50] Antistress activity ^[51] Memory-enhancing effects ^[51] Antibacterial activity ^[52] Hepatoprotective activity ^[53] Diuretic ^[54] Anti-osteoporotic activity ^[55] Hypolipidemic Activity ^[56] Emmenagogue and galactogogue, estrogenic activity ^[57]

		(bergapten) and Sterols (β -sitosterol) [16,18,46,47]	<i>Kāsir-i-Riyah</i> (carminative), <i>Muḥallil</i> (resolvent), <i>Mudirr-i-Laban</i> (galactogogue), <i>Musakkin-i-Alam</i> (analgesic) [13-15,33,47]	
4.	<i>Gaozaban</i> Arabic: <i>Lisan-us-Saur</i> English: Borage [18, 59]	Pyrrolizidine alkaloids, licosamin, intermedin, sopinin, sopindian, fatty acids (ALA & GLA), silicic acid, potassium, calcium, nitrate acetic, lactic and malic acid, δ -bornesitol, cianozhens, mucilage. [18,58,59]	<i>Dāfi'-i- Ḥummā</i> (anti-pyretic), <i>Muqawwī</i> (tonic), <i>Muddir-i-Bawl</i> (diuretic), <i>Muqawwī-i-Qalb</i> (cardiotonic), <i>Mufarriḥ</i> (exhilarant), <i>Mulaiyyin</i> (laxative), <i>Munaffith</i> (expectorant), <i>Mufattit-i-Ḥaṣāh</i> (lithotriptic) [12,14]	Hepatoprotective activity [60] Cardiovascular activity [61] Respiratory activity [61] Gastrointestinal activity [61] Antioxidant activity [62] Antinociceptive activity [63,65] Anxiolytic activity [64] Anti-inflammatory activity [65] Antibacterial activity [66]
5.	Maweez munaqqa Arabic: Zabeeb-ul-Jabal English: Raisins [16,15,18]	Flavonoids (kaempferol-3-O-glucosides, quercetin-3-O-glucosides, quercetin and myricetin) Polyphenols (flavan-3-ol derivatives) Catechins, Epicatechin, Epicatechin-3-O-gallate, Procyanidins or proanthocyanidins [16,18, 67]	<i>Mughadhdhī</i> (Nutritive), <i>Mundiz-e-Khilṭ ghaleez</i> (concoctive of viscous humour), <i>Mufattiḥ -i-Sudud</i> (deobstruent), <i>Mulaiyyin-i-Shikam</i> (laxative), <i>Muḥallil</i> (resolvent), <i>Jālī</i> (degergent), <i>Muqawwī-i-Mi'da wa Am'ā</i> (tonic for stomach and intestine), <i>Muqawwī-i-Jigar</i> (hepatotonic), <i>Muḥarriq-e-Bāh</i> (stimulant), <i>Musammin-i-Badan</i> (adipogenous) [12-14,68]	Hepatoprotective activity [69] Antioxidant activity [70] antidiabetic activity [70] Antiulcer activity [36] Antioxidant effect [71] Anti-hypercholesterolemic potential [71] Cardioprotective activity [72] Antitumour activity [73] Antibacterial activity [74] Anti-stress activity [75] Antifungal activity [76] Anti-inflammatory activity [77]

DISCUSSION

Nuskha-e-khalal-e-shikam is very efficacious medicine, the prescription of *NKS* is mentioned in various classical texts. Various functions of *Nuskha-e-khalal-e-shikam* is attributed to its various ingredients which possess antipyretic, antioxidant, hepatoprotective activity, antibacterial, expectorant, anti-inflammatory, antiviral effect, hypolipidemic activity, carminative, tonic for nerves, brain, heart and kidneys, nerve stimulant, immunomodulator activity, emmenagogue, laxative, deobstruent, demulcent, blood purifier, anxiolytic activity, analgesic etc properties. These functions have been proved by scientific studies.

Most of the ingredients of the *NKS* contains flavonoids, terpenes and Coumarins.

Flavonoids

As studies shown that flavonoids have capacity to act as antioxidants. The flavones and catechins seem to be the most powerful flavonoids for protecting the body against reactive oxygen species. flavonoids are well known as antibacterial agents against a wide range of pathogenic microorganism [78]. Anti-inflammatory and antipyretic feature are the key abilities of flavonoids. [78][79]

Essential oil

V. odorata, *F. vulgare* and *C. intybus* contain essential oils that include terpenoids, phenylpropanoids. *C. intybus* is particularly rich in a wide range of natural terpenoids named sesquiterpene lactones. Studies shows that sesquiterpene lactones have antibacterial antiviral anti-inflammatory activities.^[80]

Coumarins

C. intybus and *F. vulgare* in the NSK contain coumarins and its derivatives. Coumarins and their derivatives exert a vast array of bioactive properties such as anticoagulant, antibacterial, anti-inflammatory, antioxidant, antitumor, antiviral.^[81]

In Unani medicine, many diseases are indeed believed to originate from *Su-i-Hazm* (faulty or incomplete digestion), because disturbed digestion produces abnormal humors (*Akhlāt*) that then affect the whole body. This concept can be correlated with modern ideas linking chronic dyspepsia, malabsorption, gut dysbiosis, low-grade inflammation, and metabolic disturbance to systemic diseases.^[82] Gut dysbiosis is a disruption of normal gut microbial balance, involving loss of beneficial microbes, overgrowth of potentially harmful ones, or reduced overall microbial diversity.

The gut microbiota, a complex ecosystem of microorganisms, regulates immune responses and maintains intestinal homeostasis. Gut dysbiosis, or damaged gut microbioma, contributes to inflammatory diseases like IBD (ulcerative colitis/Crohn's), rheumatoid arthritis, lupus, and skin issues (psoriasis, eczema) by boosting harmful bacteria like *Prevotella* and reducing helpful ones like *Faecalibacterium*. This triggers excess inflammation via cytokines like IL-17. It also worsens infections such as COVID-19 and TB through low short-chain fatty acids and bacterial shifts.^[83]

Phytochemicals such as polyphenols, flavonoids, terpenoids, saponins, and capsaicin play a pivotal role in modulating the gut microbiota by suppressing pathogenic bacteria and promoting the growth of beneficial genera.^[84] The active chemical constituents present in the ingredients of *Nuskha-e-Khalal-e-Shikam*, including flavonoids, polyphenols, coumarins, terpenes, terpenoids, and saponins, exert potent antioxidant and anti-inflammatory effects, which help to repair and to restore the gut microbiome. Consequently, these actions enhance intestinal barrier integrity, strengthen immune function, and contribute to the prevention of inflammatory and metabolic disorders.

However, clinical trials or scientific studies are lacking on *Nuskha-e-khalal-e-shikam* as a compound drug.

CONCLUSION

With the above discussion the inference may be drawn that the *Nuskha-e-khalal-e-shikam* is one of the best Unani formulations with a lot of health benefits. *Nuskha-e-khalal-e-shikam* is effective in the management of several undiagnosed condition of the patient since time immemorial, so it has been found to be a time-tested drug. However more scientific studies and clinical trials are needed on this compound formulation to ensure its scientific validation for clinical use in patients.

REFERENCES

- Ahmad S. I. Kulliyate Asri, New public press Delhi; 4-12, 1983.
- Nafees B.: Kulliyat-e-Nafeesi (Tarjuma wa sharah by M. Kabiruddin), Idara Kitab-ul-shifa, Daryaganj, New Delhi 1954; pp 16-81, 517-520, 453.
- Hamdani K.H, Usool-e-Tib, Qaumi council Bararye Farogh Urdu Zuban, New Delhi, 2011, 39-61, 405, 431.
- Ibn Sina: Kulliyat-e-Qanoon (Tarjuma wa sharah by M. Kabiruddin), Aijaz Publishing House, New Delhi 2006; pp 253-255.
- Husain, A., Sofi, G.D., Tajuddin, T., Dang, R. and Kumar, N., 2010. Unani system of medicine-introduction and challenges. Medical Journal of Islamic World Academy of Sciences, 18(1), pp.27-30.
- Kejariwal, D., Sarkar, N., Chakraborti, S.K., Agarwal, V. and Roy, S., 2001. Pyrexia of unknown origin: a prospective study of 100 cases. Journal of postgraduate medicine, 47(2), p.104.
- Petersdorf RG, Beeson PB. Fever of unexplained origin: Report on 100 cases. Medicine (Baltimore) 1961; 40:1-30.
- Doyle, J.F. and Schortgen, F., 2016. Should we treat pyrexia? And how do we do it?. Critical Care, 20(1), pp.1-10.
- Kabiruddin M. Sharah Asbab, Vol- 3&4. Idara Kitab ul shifa, New Delhi 2010; pp- 387-394, 416-420.
- Ibn Sina: Hummiyat-e-Qanoon (Urdu translation by M. Kabiruddin), Aijaz Publishing House, New Delhi 2007; pp- 227-229.
- Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet; <https://powo.science.kew.org/>. Accessed 2026 Feb 28.
- Ghani HN. Khazāin al-advīā, Vol. 1 to 4th. 3rd edition. New Delhi: Idara Kitab-ul-shifa; 2011. pp. 397-398, 997, 1116-1117, 1284

13. Khān MA. Muḥīṭ-i A'zam, (Urdu translation), First edition. New Delhi: Central Council for Research in Unani Medicine, Ministry of AYUSH, Government of India; 2012. Pp Vol-1. 771-774, 543-547; Vol-IV – 18-23, 692-695.
14. Kabiruddin HM. Makhzan-ul-Mufradat-Almaroof-Khwas-ul-adwia. Faisal Publications Deoband 2000; pp- 138, 428-429, 112-113, 481-482, 560-561.
15. Anonymous, The Unani Pharmacopoeia of India, Govt. of India, Ministry of Health & Family Welfare, Dept. of AYUSH, New Delhi, Part-1, Vol. IV, pp- 96-97, Part-1 Vol. III- 14-16, Part-1 Vol-1- 15-16
16. Nadkarni KM. Indian Materia Medica, Vol-1. 3rd Edition. Mumbai: popular Prakashan; 1976. Pp- 1274-1275, 313-314, 557-559, 1285-1289.
17. Anonymous. Standardization of Single Drugs of Unani Medicine. Part II. New Delhi: Central Council for Research in Unani Medicine (CCRUM). Ministry of Health & Family Welfare, Govt. of India, 1992; 149- 152.
18. Khare CP. Indian Medicinal Plants. First edition. New Delhi: Springer publication; 2007. pp. 706, 146-147, 271, 97, 711-712
19. Fazeenah A. and Quamri M.A. 2020. Banafsha (*Viola Odorata* Linn.) -A Review. World Journal of Pharmaceutical Research, 9(10): 514-537.
20. Gautam S.S. and Kumar S., 2012. The antibacterial and phytochemical aspects of *Viola odorata* Linn. extracts against respiratory tract pathogens. Proceedings of the National Academy of Sciences, India Section B: Biological Sciences, 82(4), pp.567-572.
21. Koochek, M.H., Pipelzadeh, M.H. and Mardani, H., 2003. The effectiveness of *Viola odorata* in the prevention and treatment of formalin-induced lung damage in the rat. Journal of herbs, spices & medicinal plants, 10(2), pp.95-103.
22. Ibraheem, R.M., Mhawesh, A.A. and Abood, K.W., 2018. Estimation of the whole flavonoid, antioxidant, antibacterial challenge concerning *viola odorata* (banafsha) methanolic extract. The Iraqi Journal of Agricultural Science, 49(4), p.655.
23. Mohammed, R., 2017. Immunological, Cytogenetic and Hepatoprotective Effect of *Viola odorata* Methanolic Extract on Methotrexate Induced Albino Male Mice. Journal of Biotechnology Research Center, 11(2), pp.47-55.
24. Asheesh, K., Suresh, C. and Meenakshi, P., 2017. A brief knowledge of banafsha (*Viola odorata* linn.) & other *Viola* species. International Journal of Ayurveda and Pharma Research.
25. Vishal, A., Parveen, K., Pooja, S., Kannappan, N. and Kumar, S., 2009. Diuretic, laxative and toxicity Studies of *Viola odorata* aerial parts. Pharmacol online, 1, pp.739-748.
26. Khattak, S.G., Gilani, S.N. and Ikram, M., 1985. Antipyretic studies on some indigenous Pakistani medicinal plants. Journal of ethnopharmacology, 14(1), pp.45-51.
27. Siddiqi, H.S., Mehmood, M.H., Rehman, N.U. and Gilani, A.H., 2012. Studies on the antihypertensive and antidyslipidemic activities of *Viola odorata* leaves extract. Lipids in health and disease, 11(1), pp.1-12.
28. Hassan, F. and Naeem, I., 2014. Biological activity of *Viola odorata* Linn against *Mycobacterium tuberculosis*. International Journal of Pharmacy and Biological Sciences, 5(3), pp.61-69.
29. Feyzabadi, Z., Jafari, F., Kamali, S.H., Ashayeri, H., Aval, S.B., Esfahani, M.M. and Sadeghpour, O., 2014. Efficacy of *Viola odorata* in treatment of chronic insomnia. Iranian Red Crescent Medical Journal, 16(12).
30. Katoch, M., Paul, A., Singh, G. and Sridhar, S.N.C., 2017. Fungal endophytes associated with *Viola odorata* Linn. as bioresource for pancreatic lipase inhibitors. BMC complementary and alternative medicine, 17(1), pp.1-8.
31. Zaman, R. and Basar, S.N., 2013. A review article of Beekhe Kasni (*Cichorium intybus*) its traditional uses and pharmacological actions. Res J Pharm Sci, 2(8), 1-4.
32. Saxena, R.A.H.U.L., Sulakhiya, K.B. and Rathore, M.A.N.O.J., 2014. *Cichorium intybus* Linn.: a review of pharmacological profile. International Journal of Current Pharmaceutical Research, 6(4), pp.11-15.
33. Hakeem M. A, Bustan-ul Mufaradat Jadeed. Idara Kitab-ush-shifa 2002. Pp-414-415, 356-357.
34. Khalid, A., Shahid, S., Khan, S.A., Kanwal, S., Yaqoob, A., Rasool, Z.G. and Rizwan, K., 2018. Antioxidant activity and hepatoprotective effect of *Cichorium intybus* (Kasni) seed extract against carbon tetrachloride-induced liver toxicity in rats. Tropical Journal of Pharmaceutical Research, 17(8), pp.1531-1538.
35. Ali, A., Jehangir, A. and Ayub, F., 2016. Hepatoprotective effect of aqueous extract of *Cichorium intybus* roots on isoniazid induced hepatotoxicity. JIIMC, 11(3), pp.99-102.
36. Pasha, S., Mahurkar, N. and Jayaveera, K.N., 2016. Protective effect of *Vitis vinifera* and *Cichorium intybus* on adrenocortical activity in stress induced albino rats. Research Journal of Pharmacy and Technology, 9(5), pp. 521-526.

37. Saxena, R., Belemkar, S., Apte, K.G., Saxena, R.C. and Bharti, M., 2011. Antiulcerogenic and Antioxidant Activity of Hydro alcoholic extract of the root of *Cichorium intybus* L. in experimentally induced ulcer in Rats. *Ethnopharmacology*, 2, pp.1-4.
38. Nandagopal, S. and Kumari, B.R., 2007. Phytochemical and antibacterial studies of Chicory (*Cichorium intybus* L.)-A multipurpose medicinal plant. *Advances in Biological Research*, 1(1-2), pp.17-21.
39. Ghamarian, A., Abdollahi, M., Su, X., Amiri, A., Ahadi, A. and Nowrouzi, A., 2012. Effect of chicory seed extract on glucose tolerance test (GTT) and metabolic profile in early and late-stage diabetic rats. *DARU Journal of Pharmaceutical Sciences*, 20(1), pp.1-9.
40. Eslami, H., Babaei, H., Falsafi, P., Rahbar, M., Najarkarimi, F. and Pourzare-Mehrbani, S., 2017. Evaluation of the Antifungal Effect of Chicory Extracts on *Candida Glabrata* and *Candida Krusei* in a Laboratory Environment. *The Journal of Contemporary Dental Practice*, 18(11), pp.1014-1020.
41. Rahimipour, A., Dehghan Nayeri, N., Mehrandish, R. and Awsat Mellati, A., 2017. Anti-Cancer Activity of Methanol Extracts of *Cichorium Intybus* on Human Breast Cancer SKBR3 Cell Line. *Razavi International Journal of Medicine*, 5(1).
42. Abdel-Rahman, R.F., Soliman, G.A., Yusufoglu, H.S., Tatli-Çankaya, I., Alqasoumi, S.I., Anul, S.A. and Akaydin, G., 2015. Potential anticonvulsant activity of ethanol extracts of *Cichorium intybus* and *Taraxacum serotinum* in rats. *Tropical Journal of Pharmaceutical Research*, 14(10), pp.1829-1835.
43. Amirghofran, Z., Azadbakht, M. and Karimi, M.H., 2000. Evaluation of the immunomodulatory effects of five herbal plants. *Journal of ethnopharmacology*, 72(1-2), pp.167-172.
44. Wesołowska, A., Nikiforuk, A., Michalska, K., Kisiel, W. and Chojnacka-Wójcik, E., 2006. Analgesic and sedative activities of lactucin and some lactucin-like guaianolides in mice. *Journal of ethnopharmacology*, 107(2), pp.254-258.
45. Nayeemunnisa and Rani, M.K., 2003. Cardioprotective effects of *Cichorium intybus* in ageing myocardium of albino rats. *Current Science*, pp.941-943.
46. Jamwal, N.S., Kumar, S. and Rana, A.C., 2013. Phytochemical and Pharmacological Review on *Foeniculum Vulgare*. *Pharma Science Monitor*, 4(3).
47. Al-Snafi, A.E., 2018. The chemical constituents and pharmacological effects of *Foeniculum vulgare*-A review. *IOSR Journal of Pharmacy*, 8(5), pp.81-96.
48. Khan, N.T., 2017. Antifungal potency of *Foeniculum vulgare* seed extract. *J. Tissue Sci. Engin*, 8(3), p.1000207.
49. Choi, E.M. and Hwang, J.K., 2004. Anti-inflammatory, analgesic and antioxidant activities of the fruit of *Foeniculum vulgare*. *Fitoterapia*, 75(6), pp.557-565.
50. Abou El-Soud, N., El-Laithy, N., El-Saeed, G., Wahby, M., Khalil, M., Morsy, F. and Shaffie, N., 2011. Antidiabetic activities of *Foeniculum vulgare* Mill. essential oil in streptozotocin-induced diabetic rats. *Macedonian J Med Sci*, 4(2), pp.139-146.
51. Koppula, S. and Kumar, H., 2013. *Foeniculum vulgare* Mill (Umbelliferae) attenuates stress and improves memory in wister rats. *Tropical Journal of Pharmaceutical Research*, 12(4), pp.553-558.
52. Khan, N., 2017. In vitro antibacterial activity of *Foeniculum vulgare* seed extract. *Agrotechnology*, 6(162), p.2.
53. Ozbek, H., 2006. Hepatoprotective effect of Fennel essential oil. *Research Journal of Medicine and Medical Science*, 1(2), pp.72-76.
54. Tanira, M.O.M., Shah, A.H., Mohsin, A., Ageel, A.M. and Qureshi, S., 1996. Pharmacological and toxicological investigations on *Foeniculum vulgare* dried fruit extract in experimental animals. *Phytotherapy Research*, 10(1), pp.33-36.
55. Fariba, J., Alireza, G. and Hossein, N., 2006. Evaluation of Prophylactic effect of the Fennel essential oil on experimental osteoporosis models in Rats. *Int J Pharmacol*, 2, pp.588-592.
56. Garg, C., Ansari, S.H., Khan, S.A. and Garg, M., 2011. Effect of *Foeniculum vulgare* Mill. fruits in obesity and associated cardiovascular disorders demonstrated in high fat diet fed albino rats. *J Pharm Biomed Sci*, 8(8), pp.1-5.
57. Mallni, T., Vanithakumari, G., Devi, N.M.S.A.K. and Fiango, V., 1985. Effect of foeniculuai vulgare. Mill seed extract on the genital organs of male and female rats. *Indian J Physiol Pharmacol*, 29, pp.21-26.
58. Asadi-Samani, M., Bahmani, M. and Rafieian-Kopaei, M., 2014. The chemical composition, botanical characteristic and biological activities of *Borago officinalis*: a review. *Asian Pacific journal of tropical medicine*, 7, pp. S22-S28.
59. Nyeem, M.A.B., Haque, M.S. and Hoque, M.A., 2017. Phytoconstituents and pharmacological activity of *Gauzaban* (*Borago officinalis* Linn): A review. *International Journal of Food Science and Nutrition*, 2(1), pp.148-152.
60. Hamed, A.N. and Wahid, A., 2015. Hepatoprotective activity of *Borago officinalis* extract against CCl4-

- induced hepatotoxicity in rats. *J. of Nat. Prod*, 8, pp.113-122.
61. Gilani, A.H., Bashir, S. and Khan, A.U., 2007. Pharmacological basis for the use of *Borago officinalis* in gastrointestinal, respiratory and cardiovascular disorders. *Journal of Ethnopharmacology*, 114(3), pp. 393-399.
 62. Mhamdi, B., Wannes, W.A., Sriti, J., Jellali, I., Ksouri, R. and Marzouk, B., 2010. Effect of harvesting time on phenolic compounds and antiradical scavenging activity of *Borago officinalis* seed extracts. *Industrial Crops and Products*, 31(1), pp. e1-e4.
 63. Shahraki, M.R., Ahmadimoghadm, M. and Shahraki, A.R., 2015. The antinociceptive effects of hydroalcoholic extract of *Borago officinalis* flower in male rats using formalin test. *Basic and clinical neuroscience*, 6(4), p.285.
 64. Komaki, A., Rasouli, B. and Shahidi, S., 2015. Anxiolytic effect of *Borago officinalis* (Boraginaceae) extract in male rats. *Avicenna Journal of Neuro Psycho Physiology*, 2(1), pp.34-38.
 65. Gihan F Asaad, Alaa Q Redai, Alaa O Hakami, Fatimah Y Ghazwani, Yousra Nomier and Saeed Alshahrani (2020) "Potential Analgesic And Anti-Inflammatory Effect Of *Cuminum Cyminum* And *Borago Officinalis* In Rats And Mice", *Asian Journal of Pharmaceutical and Clinical Research*, 13(1), pp. 216–218.
 66. Karimi, E., Oskoueian, E., Karimi, A., Noura, R. and Ebrahimi, M., 2018. *Borago officinalis* L. flower: a comprehensive study on bioactive compounds and its health-promoting properties. *Journal of Food Measurement and Characterization*, 12(2), pp.826-838.
 67. Nassiri-Asl, M. and Hosseinzadeh, H., 2016. Review of the pharmacological effects of *Vitis vinifera* (Grape) and its bioactive constituents: an update. *Phytotherapy Research*, 30(9), pp.1392-1403.
 68. Khan, T., Farooqui, A.H., Ahmed, M.N. and Fatima, A., Sharbat-e-maweez is effective in soo-ul-qiniya bawajah qillat-e-faulad (iron deficiency anemia): A review. *Int J Unani Integ Med* 2017;1(2):07-09. DOI: 10.33545/2616454X.2017.v1.i2a.11
 69. Ahmad, F. and Khan, G.M., 2012. Study of aging and hepatoprotective activity of *Vitis Vinifera* L. seeds in albino rats. *Asian Pacific Journal of Tropical Biomedicine*, 2(3), pp. S1770-S1774.
 70. Şendoğdu, N., Aslan, M., Orhan, D.D., Ergun, F. and Yeşilada, E., 2006. Antidiabetic and antioxidant effects of *Vitis vinifera* L. leaves in streptozotocin-diabetic rats. *Turkish J Pharm Sci*, 3(1), pp.7-18.
 71. Devi, S. and Singh, R., 2017. Antioxidant and anti-hypercholesterolemic potential of *Vitis vinifera* leaves. *Pharmacognosy Journal*, 9(4).
 72. Cheng, M., Gao, H.Q., Xu, L., Li, B.Y., Zhang, H. and Li, X.H., 2007. Cardioprotective effects of grape seed proanthocyanidins extracts in streptozocin induced diabetic rats. *Journal of cardiovascular pharmacology*, 50(5), pp.503-509.
 73. Zhao, J., Wang, J., Chen, Y. and Agarwal, R., 1999. Anti-tumor-promoting activity of a polyphenolic fraction isolated from grape seeds in the mouse skin two-stage initiation–promotion protocol and identification of procyanidin B5-3'-gallate as the most effective antioxidant constituent. *Carcinogenesis*, 20(9), pp.1737-1745.
 74. Jayaprakasha, G.K., Selvi, T. and Sakariah, K.K., 2003. Antibacterial and antioxidant activities of grape (*Vitis vinifera*) seed extracts. *Food research international*, 36(2), pp.117-122.
 75. Pasha, S., Mahurkar, N. and Jayaveera, K.N., 2014. Evaluation of Antistress Activity of *Vitis vinifera* and *Chicorium intybus*. *Research Journal of Pharmacy and Technology*, 7(12), pp.1377-1381.
 76. Fraternali, D., Ricci, D., Verardo, G., Gorassini, A., Stocchi, V. and Sestili, P., 2015. Activity of *Vitis vinifera* tendrils extract against phytopathogenic fungi. *Natural product communications*, 10(6), p.1934578X1501000661.
 77. Di Lorenzo, C., Sangiovanni, E., Fumagalli, M., Colombo, E., Frigerio, G., Colombo, F., Peres de Sousa, L., Altindişli, A., Restani, P. and Dell'Agli, M., 2016. Evaluation of the anti-inflammatory activity of raisins (*Vitis vinifera* L.) in human gastric epithelial cells: a comparative study. *International journal of molecular sciences*, 17(7), p.1156.
 78. Xie Y, Yang W, Tang F, Chen X, Ren L. Antibacterial activities of flavonoids: structure-activity relationship and mechanism. *Current medicinal chemistry*. 2015 Jan 1; 22(1): 132-49.
 79. Khattak SG, Gilani SN, Ikram M. Antipyretic studies on some indigenous Pakistani medicinal plants. *Journal of ethnopharmacology*. 1985 Sep 1; 14(1): 45-5
 80. Sülsen VP, Martino VS. Sesquiterpene lactones. *Advances in their Chemistry and Biological Aspects*. Cham: Springer. 2018: 379.
 81. Garg SS, Gupta J, Sharma S, Sahu D. An insight into the therapeutic applications of coumarin compounds and their mechanisms of action. *European Journal of Pharmaceutical Sciences*. 2020 Sep 1; 152: 105424.
 82. Ul Haq E, Kumar R. Understanding the process of Hazm (digestion) in Unani medicine: a

- physiological correlation with modern GI function. International Journal of Unani and Integrative Medicine. 2025; 9(2): 190-192. doi: 10.33545/2616454X.2025.v9.i2c.359.
83. Maciel-Fiuza MF, Muller GC, Campos DM, do Socorro Silva Costa P, Peruzzo J, Bonamigo RR, Veit T, Vianna FS. Role of gut microbiota in infectious and inflammatory diseases. Frontiers in microbiology. 2023 Mar 27; 14: 1098386.
84. Singh A, Kaur P, Kumar M, Shafi S, Upadhyay PK, Tiwari A, Tiwari V, Rangra NK, Thirunavukkarasu V, Kumari S, Roy D. The Role of Phytochemicals in Modulating the Gut Microbiota: Implications for Health and Disease. Medicine in Microecology. 2025 Mar 27: 100125.

Cite this article as:

Uzma Viquar, Sadaf Joweria, Mubasshera Mohd. Irfan, Mohd. Noman Taaha, Younis Iftikhar Munshi. Therapeutic Efficacy and Phytochemical Insights into the Classical Unani Formulation Nuskha-E-Khalal-E-Shikam. AYUSHDHARA, 2026;13(2):49-58.

<https://doi.org/10.47070/ayushdhara.v13i2.2557>

Source of support: Nil, Conflict of interest: None Declared

***Address for correspondence**

Dr. Uzma Viquar

Professor

PG Dept. of Ilmu Advia (unani pharmacology) National Research Institute of Unani Medicine for Skin Disorder, Hyderabad, Telangana.

Email: viquar.uzma@gmail.com

Disclaimer: AYUSHDHARA is solely owned by Mahadev Publications - A non-profit publications, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. AYUSHDHARA cannot accept any responsibility or liability for the articles content which are published. The views expressed in articles by our contributing authors are not necessarily those of AYUSHDHARA editor or editorial board members.

