



Review Article

HOMOEOPATHIC APPROACH IN THE MANAGEMENT OF PILONIDAL SINUS

Mayuri Suryawanshi^{1*}, Sundara Jyothi Bhagvatsingh²

¹PG Scholar, ²Guide, Department of Repertory, Guru Mishri Homoeopathic Medical College and Hospital, Shelgaon, Maharashtra, India.

Article info

Article History:

Received: 11-07-2026

Accepted: 07-08-2026

Published: 10-09-2026

KEYWORDS:

Pilonidal sinus,
Pilonidal disease,
Natal cleft,
Homoeopathy,
BBCR, Repertory.

ABSTRACT

Pilonidal sinus disease (PSD) is a chronic disorder of the natal cleft that predominantly affects adolescents and young adults. It may present as an acute abscess or as a chronic discharging sinus, with symptoms including pain, swelling, local tenderness and intermittent discharge. The contemporary understanding of pilonidal disease supports an acquired pathogenesis associated with hair penetration, follicular obstruction and local mechanical factors. Management depends upon the clinical presentation and may range from local care and observation to incision and drainage or definitive surgical procedures. Homoeopathy approaches patients on the basis of individual symptom patterns and traditionally employs materia-medica and repertorial methods for remedy selection. In this article, the application of Boger Boenninghausen's Characteristics and Repertory (BBCR) to symptoms associated with pilonidal sinus is discussed. Selected remedies traditionally considered in suppurative and chronic sinus conditions, including *Silicea*, *Hepar sulphuris calcareum*, *Myristica sebifera* and *Calcarea sulphurica*, are described from the homoeopathic materia-medica perspective. The article primarily presents a homoeopathic and repertorial perspective and should not be interpreted as evidence that homoeopathic treatment can replace established surgical management of acute abscesses or complicated pilonidal disease. Further well-designed clinical research is required to determine the effectiveness and safety of homoeopathic interventions in this condition.

INTRODUCTION

Pilonidal sinus disease is a condition involving the skin and subcutaneous tissue of the natal cleft. It occurs predominantly in younger individuals and may range from asymptomatic pits to painful abscesses and chronically discharging sinus tracts. Current evidence considers pilonidal disease to be primarily acquired rather than a congenital abnormality. Hair penetration into the natal cleft and subsequent foreign-body inflammation are important components of its pathogenesis. [1-4]

Several factors have been associated with the development of pilonidal disease, including prolonged sitting, local friction or trauma, obesity, a deep natal

cleft, excessive hair and a positive family history. The disease may significantly interfere with sitting, walking, work and other daily activities, particularly when recurrent inflammation or chronic discharge occurs. [1,2]

Treatment is determined by the stage and severity of disease. An acute pilonidal abscess generally requires appropriate surgical drainage, whereas chronic or recurrent disease may require definitive procedures. A variety of minimally invasive and surgical techniques have been described, and recurrence remains an important consideration when treatment options are compared. [2-5]

From the homoeopathic perspective, the individual symptom picture is considered during remedy selection. Repertorial analysis may assist in organizing characteristic symptoms according to location, sensation, modalities, concomitants and general symptoms. The present article discusses the application of BBCR in the analysis of symptoms occurring in pilonidal sinus disease.

Access this article online

Quick Response Code



<https://doi.org/10.47070/ayushdhara.v13i4.2947>

Published by Mahadev Publications (Regd.)
publication licensed under a Creative Commons
Attribution-NonCommercial-ShareAlike 4.0
International (CC BY-NC-SA 4.0)

Definition

Pilonidal disease is an acquired inflammatory condition affecting the natal cleft, in which hair and keratinous material may become associated with pits, cystic cavities or sinus tracts. The disease may remain asymptomatic or progress to an acute abscess or chronic discharging sinus. ^[1-4]

The characteristic clinical finding is one or more pits or openings in the midline of the natal cleft. Chronic disease may be associated with lateral secondary openings, recurrent inflammation and persistent discharge. ^[1]

Epidemiology

Pilonidal disease is predominantly a disorder of adolescents and young adults. Published studies indicate a male predominance, although females are also affected. The reported incidence varies considerably between populations and studies. ^[1,2]

Important factors associated with pilonidal disease include:

- Prolonged sitting or sedentary occupation
- Increased local hair density
- Obesity
- Deep natal cleft
- Local friction or trauma
- Sweating and local moisture
- Family history
- Previous episodes of pilonidal disease. ^[1-4]

Because these factors may vary between individuals, assessment of the patient's occupation, body habitus, local hygiene and previous episodes is relevant during clinical evaluation.

Etiology and Pathogenesis

The currently accepted understanding of pilonidal disease favours an acquired mechanism. Loose hairs may become trapped within the natal cleft and penetrate the skin. Alternatively, follicular occlusion may initiate inflammation and subsequent sinus formation. The embedded hair acts as a foreign material and produces a local inflammatory response. ^[1,2]

Mechanical forces, friction and the anatomy of the natal cleft may facilitate hair penetration and retention. Repeated inflammation can result in abscess formation and the development of chronic sinus tracts. Histological studies have demonstrated inflammatory tissue and foreign-body reactions surrounding embedded hair. ^[1]

The pathogenesis may therefore be represented as:

Hair/follicular obstruction → local inflammation → pit formation → abscess formation → drainage → chronic sinus tract formation

This process may be modified by local hygiene, hair characteristics, sweating, friction and individual anatomical factors. ^[1-4]

Pathology

The pathological process generally begins with obstruction or penetration involving hair follicles in the natal cleft. Local inflammation develops around the retained hair or keratinous material. Continued inflammation can result in an abscess cavity, which may subsequently communicate with the skin through sinus openings. ^[1,2]

Repeated episodes may produce chronic granulation tissue, fibrosis and multiple sinus tracts. The extent of disease can therefore vary from a small solitary pit to complex recurrent disease with several interconnected tracts. ^[1,4]

Clinical Features

Clinical presentation depends upon whether the disease is asymptomatic, acute, chronic or recurrent.

Acute Pilonidal Abscess

Typical findings include:

- Sudden or progressively increasing pain
- Localized swelling in the natal cleft
- Erythema and warmth
- Tenderness
- Fluctuant abscess formation
- Purulent discharge following spontaneous or surgical drainage.

Systemic symptoms such as fever or malaise may occur when significant infection is present. ^[1,3]

Chronic Pilonidal Disease

Chronic disease may present with:

- One or more pits in the natal cleft
- Intermittent or persistent discharge
- Recurrent discomfort or pain
- Local irritation
- Granulation tissue
- Secondary sinus openings.

The discharge may be serous, purulent or occasionally blood-stained. ^[1,2]

Recurrent or Complex Disease

Recurrent disease may involve several sinus openings, lateral extension of tracts, recurrent abscess formation and fibrosis. Complex disease may require more extensive assessment and treatment. ^[1,4]

Diagnosis

Diagnosis is principally clinical. A careful history and examination of the natal cleft are usually sufficient in uncomplicated cases. ^[1,3]

Clinical Examination

The examination should assess:

- Number and location of pits
- Presence of discharge
- Tenderness and swelling
- Erythema or local inflammation
- Induration
- Abscess formation
- Lateral extension of sinus openings.

Routine imaging is not required in most uncomplicated cases. Imaging may be considered when the diagnosis is uncertain or when recurrent or complex disease requires further anatomical assessment. [1,3]

Differential Diagnosis

Conditions that may resemble pilonidal disease include:

1. Perianal abscess– Generally located close to the anus and associated with perianal pain or swelling.
2. Furuncle– Usually a localized follicular infection rather than a chronic sinus tract.
3. Hidradenitis suppurativa– Recurrent inflammatory nodules and sinus formation involving characteristic intertriginous areas.
4. Anal fistula– Typically associated with an opening close to the anal canal and a history suggestive of anorectal disease.
5. Other chronic sinuses or inflammatory lesions– May occasionally require further investigation to establish the diagnosis.

Correct differentiation is important because management varies substantially between these conditions.

Conventional Management: Brief Overview

Management should be based on the clinical presentation.

An acute pilonidal abscess is generally managed by incision and drainage rather than attempting definitive treatment of the chronic sinus during the acute episode. Chronic or recurrent disease may require procedures designed to eliminate the sinus tracts and reduce recurrence. [3,4]

Several operative techniques have been described. Evidence from systematic reviews indicates that, when surgical closure is performed, off-midline closure generally has advantages over conventional midline closure with respect to healing and recurrence.[5]

Non-operative measures such as attention to natal-cleft hygiene and management of hair may also form part of preventive care, although the evidence for individual preventive interventions varies. [6]

Homoeopathic Perspective

In classical homoeopathic practice, remedy selection is based on the totality of symptoms rather than solely on the diagnostic label. Features such as the character of pain, nature of discharge, local sensitivity, modalities, concomitant symptoms and general characteristics may be considered during case analysis.

In the context of pilonidal sinus, remedies traditionally associated with suppuration, abscess formation and chronic sinus conditions may be considered according to the individual symptom picture. The following descriptions represent traditional materia-medica indications and should not be interpreted as evidence of proven clinical efficacy.

1. *Silicea*

Traditionally considered in chronic suppurative conditions, particularly when there is a tendency toward recurrent abscess formation, delayed healing and persistent discharge. In homoeopathic materia medica, *Silicea* is also associated with conditions involving retained foreign material and chronic fistulous processes.

2. *Hepar sulphuris calcareum*

Traditionally considered when an inflammatory or suppurative lesion is highly sensitive and painful. The remedy picture is commonly associated with marked tenderness, abscess formation and thick purulent discharge.

3. *Myristica sebifera*

Traditionally described in homoeopathic literature in relation to abscesses and suppurative conditions. It has historically been referred to as a remedy for facilitating suppuration and drainage.

4. *Calcarea sulphurica*

Traditionally considered in chronic suppurative conditions with persistent discharge and delayed healing. Its materia-medica picture includes prolonged suppuration and difficulty in complete healing.

5. *Fluoricum acidum*

Traditionally considered in chronic conditions associated with induration and destructive or ulcerative tendencies. Its selection would depend upon the totality rather than the diagnosis alone.

6. *Mercurius solubilis*

Traditionally associated with inflammatory and suppurative conditions accompanied by offensive secretions and characteristic systemic symptoms.

7. *Lycopodium clavatum*

The traditional materia-medica picture includes chronic suppurative states together with characteristic digestive and constitutional symptoms.

Remedy selection should therefore depend upon the complete symptom picture.

8. *Arnica montana*

Traditionally associated with conditions following trauma and with sensations of soreness or bruising. Its use in a pilonidal case would require appropriate characteristic symptoms rather than diagnosis alone.

9. Sulphur

Traditionally considered in recurrent skin disorders and chronic suppurative conditions, particularly when characteristic thermal and itching symptoms are present.

10. Graphites

Traditionally associated with chronic skin conditions and sluggish healing, particularly where thick, sticky or glutinous secretions are present.

BBCR Repertorial Approach

Boger's Boenninghausen's Characteristics and Repertory (BBCR) is a repertorial work developed from the Boenninghausen tradition and expanded by C. M. Boger. The original work was published in 1905, with later editions and publications appearing subsequently. [7,8]

The repertorial method emphasizes characteristic symptoms and allows symptoms to be analysed according to their location, sensation, modalities and associated features. The BBCR therefore provides a framework through which the patient's individual symptom expression can be organized before comparison with the materia medica. For pilonidal sinus cases, relevant symptom groups may include symptoms relating to the sacrum and coccyx, suppuration and the character of secretions.

Relevant BBCR Rubrics Used for Case Analysis

1. BACK – Sacrum and coccyx – Coccyx – Painful
2. BACK – Sacrum and coccyx – Pain – Simple, aching
3. BACK – Sacrum and coccyx – Itching
4. BACK – Sacrum and coccyx – Suppurative pain
5. Sensation and Complaints in General – Suppuration
6. Sensation and Complaints in General – Suppuration – Pus, bloody
7. Sensation and Complaints in General – Discharge, secretions – Acrid
8. Sensation and Complaints in General – Discharge, secretions – Albuminous
9. Sensation and Complaints in General – Discharge, secretions – Bland
10. Sensation and Complaints in General – Discharge, secretions – Bloody, blood-streaked
11. Sensation and Complaints in General – Discharge, secretions – Burning

12. Sensation and Complaints in General – Discharge, secretions – Corrosive [7,8]

The exact rubrics selected should correspond to symptoms actually elicited during individual case-taking.

Role of Individualized Case Analysis

A repertorial approach may be particularly useful when a patient presents with several local characteristics, modalities and concomitant symptoms. Rather than assigning a remedy solely because of the disease name, the practitioner may first identify the most characteristic features and then compare the repertorial result with the corresponding materia-medica picture.

The final prescription should therefore be based on the complete clinical and homoeopathic assessment. Reassessment during follow-up should include changes in pain, discharge, swelling, local inflammation and general well-being.

DISCUSSION

Pilonidal sinus is a relatively common disorder of young adults and may have a considerable effect on daily activities. Contemporary literature recognizes hair-related mechanisms, local mechanical factors and anatomical characteristics of the natal cleft as important contributors to disease development. [1-4]

The management of pilonidal disease remains dependent upon disease stage. Acute abscesses require appropriate drainage, while chronic and recurrent disease may require definitive surgical treatment. Evidence also indicates that surgical techniques avoiding midline closure can provide better outcomes than traditional midline closure in appropriate patients. [4,5]

Within homoeopathic practice, the disease may be analysed through individual symptoms and repertorial methods. BBCR provides a historical repertorial framework for organizing characteristic symptoms and comparing corresponding remedies. [7,8]

However, the therapeutic claims in this article should be interpreted cautiously. The traditional materia-medica descriptions of remedies should not be equated with clinical evidence demonstrating efficacy in pilonidal sinus disease. At present, the available conventional evidence base does not establish homoeopathy as a replacement for surgical treatment of acute abscesses or complex pilonidal disease. Well-designed prospective clinical studies, with appropriate diagnostic criteria, outcome measures, follow-up and comparison groups, would be necessary to evaluate any specific homoeopathic intervention.

CONCLUSION

Pilonidal sinus is a chronic or recurrent disorder of the natal cleft that predominantly affects young adults. Its development is associated with hair penetration or follicular disease together with local mechanical and anatomical factors. Diagnosis is usually clinical, and treatment should be selected according to disease severity.

From a homoeopathic perspective, individualization and repertorial analysis may be used to organize the patient's characteristic symptoms. BBCR provides a repertorial framework in which symptoms relating to location, sensation, suppuration, discharge and modalities can be evaluated.

Remedies such as *Silicea*, *Hepar sulphuris calcareum*, *Myristica sebifera* and *Calcarea sulphurica* have traditional materia-medica associations with suppurative conditions; however, these traditional indications should not be presented as established evidence of efficacy in pilonidal sinus disease.

Further systematic and adequately designed clinical research is required to determine whether individualized homoeopathic treatment has measurable benefits in patients with pilonidal sinus disease. Conventional medical and surgical care should not be delayed when acute abscess, extensive infection, recurrent disease or other complications are present.

REFERENCES

1. Hull TL, Wu J. Pilonidal disease. Surg Clin North Am. 2002; 82(6): 1169-1185.
2. Khanna A, Rombeau JL. Pilonidal disease. Clin Colon Rectal Surg. 2011; 24(1): 46-53.
3. Chintapatla S, Safarani N, Kumar S, Haboubi N. Sacrococcygeal pilonidal sinus: historical review, pathological insight and surgical options. Tech Coloproctol. 2003; 7(1): 3-8.
4. Iesalnieks I, Ommer A. The management of pilonidal sinus. Dtsch Arztebl Int. 2019; 116(1-2): 12-21.
5. Al-Khamis A, McCallum I, King PM, Bruce J. Healing by primary closure versus open healing after surgery for pilonidal sinus: systematic review and meta-analysis. BMJ. 2010; 340: c283.
6. McCallum I, King PM, Bruce J. Healing by primary versus secondary intention after surgical treatment for pilonidal sinus. Cochrane Database Syst Rev. 2007; (4): CD006213.
7. Boger CM. Boenninghausen's Characteristics and Repertory. Parkersburg, WV: Self-published; 1905.
8. Boger CM. Boenninghausen's Characteristics and Repertory. Indian ed. Bombay: Roy & Co.; 1937.
9. Da Silva JH. Pilonidal cyst: cause and treatment. Dis Colon Rectum. 2000; 43(8): 1146-1156.
10. Akinboye A, O'Keefe K, Kavalukas SL, Barbul A. Pilonidal sinus disease: review of current practice and prospects for endoscopic treatment. World J Gastrointest Surg. 2020; 12(9): 463-477

Cite this article as:

Mayuri Suryawanshi, Sundara Jyothi Bhagvatsingh. Homoeopathic Approach in the Management of Pilonidal Sinus. AYUSHDHARA, 2026;13(4):587-591.

<https://doi.org/10.47070/ayushdhara.v13i4.2947>

Source of support: Nil, Conflict of interest: None Declared

*Address for correspondence

Dr. Mayuri Suryawanshi

PG Scholar,

Department of Repertory,

Guru Mishri Homoeopathic

Medical College and Hospital,

Shelgaon, Maharashtra, India.

Email:

Mayurisuryawanshi33@gmail.com

Disclaimer: AYUSHDHARA is solely owned by Mahadev Publications - A non-profit publications, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. AYUSHDHARA cannot accept any responsibility or liability for the articles content which are published. The views expressed in articles by our contributing authors are not necessarily those of AYUSHDHARA editor or editorial board members.